arXiv:1304.4315v1 [math-ph] 16 Apr 2013

Spectrum of the Laplacian on a covering graph with pendant edges I: The one-dimensional lattice and beyond

Akito Suzuki[*]
Department of Mathematics,
Faculty of Engineering, Shinshu University,
Wakasato, Nagano 380, Japan

April 17, 2013

Contents

1 **Introduction** **2**
- 1.1 Definitions and examples . . . 2
- 1.2 Results . . . 5

2 **Characteristic equation** **11**
- 2.1 Decomposition of the transition operator . . . 11
- 2.2 Characteristic equation . . . 13
- 2.3 Lowest degree of the characteristic polynomial . . . 19

3 **Spectrum** **24**
- 3.1 Symmetry of the spectrum . . . 24
- 3.2 Spectral gap . . . 25
- 3.3 Existence of the zero eigenvalue . . . 26

4 **Band structure of the spectrum** **27**
- 4.1 Band functions . . . 27
- 4.2 Absence of nonzero eigenvalues . . . 30
- 4.3 Multiplicity . . . 30

5 **Two dimensional lattice** **33**

---

[*] e-mail: akito@shinshu-u.ac.jp

Abstract

In this paper, we examine covering graphs that are obtained from the d-dimensional integer lattice by adding pendant edges. In the case of $d = 1$, we show that the Laplacian on the graph has a spectral gap and establish a necessary and sufficient condition under which the Laplacian has no eigenvalues. In the case of $d = 2$, we show that there exists an arrangement of the pendant edges such that the Laplacian has no spectral gap.

1 Introduction

1.1 Definitions and examples

Let $G = (V(G), E(G))$ be an undirected graph, where $V(G)$ and $E(G)$ are the sets of vertices and edges, respectively. We denote an edge that connects vertices x and y as an unordered pair $\{x, y\}$ and write $x \sim y$ if $\{x, y\} \in E(G)$. For a vertex $x \in V(G)$, we denote the number of edges incident to x as $\mathrm{deg}x$, i.e., $\mathrm{deg}x = \#\{y \mid y \sim x\}$. A vertex x is called an *end vertex* if $\mathrm{deg}x = 1$, and an edge that contains an end vertex is called a *pendant edge*. In this paper, we first consider a graph satisfying the following conditions:

- (G_1) $\{(n, 0) \in \mathbb{Z} \times \{0, 1\} \mid n \in \mathbb{Z}\} \subset V(G) \subset \mathbb{Z} \times \{0, 1\}$ and a vertex $(n, 0) \in V(G)$ is connected to $(n \pm 1, 0) \in V(G)$.
- (G_2) If $(n, 1) \in V(G)$, then $(n, 1)$ is an end vertex connected to $(n, 0)$ and $\mathrm{deg}(n, 0) = 3$. If $(n, 1) \notin V(G)$, then $\mathrm{deg}(n, 0) = 2$.

Such a graph is obtained by adding pendant edges to the one-dimensional lattice $\mathbb{Z}$, in which case a vertex $n \in \mathbb{Z}$ and an end vertex connected to $n \in \mathbb{Z}$ are identified with $(n, 0) \in V(G)$ and $(n, 1) \in V(G)$, respectively. Let

$$\ell^2(V(G)) = \left\{\psi : V(G) \to \mathbb{C} \;\middle|\; \langle\psi|\psi\rangle < \infty\right\}$$

be the Hilbert space of square summable functions on $V(G)$ with the inner product:

$$\langle\psi|\phi\rangle = \sum_{x\in V(G)} \overline{\psi}(x)\phi(x)\mathrm{deg}x, \quad \psi, \phi \in \ell^2(V(G)).$$

The transition operator L_G is a bounded self-adjoint operator on $\ell^2(V(G))$ defined as

$$(L_G\psi)(x) = \frac{1}{\mathrm{deg}x} \sum_{y\sim x} \psi(y), \quad \psi \in \ell^2(V(G)).$$

We are interested in the spectrum $\sigma(-\Delta_G)$ of the (negative) Laplacian $-\Delta_G$ defined by $-\Delta_G = 1 - L_G$. From the spectral mapping theorem, we know that $\sigma(-\Delta_G) = \{1-l \mid l \in \sigma(L_G)\}$ and that $\sigma_{\mathrm{p}}(-\Delta_G) = \{1-l \mid l \in \sigma_{\mathrm{p}}(L_G)\}$, where $\sigma_{\mathrm{p}}(L)$ denotes the set of eigenvalues of the operator L. Hence, we know that L_G has a spectral gap (resp. an eigenvalue) if and only if $-\Delta_G$ has a spectral gap (resp. an eigenvalue). In this paper, we consider the spectrum $\sigma(L_G)$ of the transition operator L_G.

A trivial example of a graph G with properties (G_1) and (G_2) is the one dimensional lattice $\mathbb{Z}$, *i.e.* G has no pendant edges. It is well known and easy to show that $\sigma(L_{\mathbb{Z}}) = [-1, 1]$ and that $L_{\mathbb{Z}}$ has no eigenvalues.

Adding a single pendant edge to each vertex of the lattice $\mathbb{Z}$, we obtain a graph $G_{1,1}$ such that $V(G_{1,1}) = \mathbb{Z} \times \{0, 1\}$ (see Figure 1, where we denote a vertex identified with an integer and an end vertex by ◯ and △, respectively). $G_{1,1}$ satisfies properties (G_1) and (G_2). Simple calculations establish that

$$\sigma(L_{G_{1,1}}) = \left[-1, -\frac{1}{3}\right] \cup \left[\frac{1}{3}, 1\right]$$

and that L_G has no eigenvalues. In particular, $L_{G_{1,1}}$ has a spectral gap.



Figure 1: Graph $G_{1,1}$.

In what follows, we consider a graph $G_{2,1}$ whose pendant edges are connected to alternate vertices of $\mathbb{Z}$: $V(G_{2,1}) = \{(n, 0) \mid n \in \mathbb{Z}\} \cup \{(2n+1, 1) \mid n \in \mathbb{Z}\}$ (see Figure 2). $G_{2,1}$ satisfies (G_1) and (G_2). Simple calculations show that

$$\sigma(L_{G_{2,1}}) = \left[-1, -\frac{1}{\sqrt{3}}\right] \cup \{0\} \cup \left[\frac{1}{\sqrt{3}}, 1\right]$$

and that 0 is an eigenvalue of $L_{G_{2,1}}$. Hence, $L_{G_{2,1}}$ has a spectral gap and a zero eigenvalue.



Figure 2: Graph $G_{2,1}$.

We note that graphs G such as $G_{1,0} \equiv \mathbb{Z}$, $G_{1,1}$ and $G_{2,1}$ have the following periodicity:

(G_3) There exists a number $r \in \mathbb{N}$ such that

$$(n+r,1) \in V(G) \text{ if and only if } (n,1) \in V(G). \tag{1.1}$$

We use $\mathscr{G}$ to denote the set of graphs that satisfy conditions (G_1) - (G_3). Each graph $G \in \mathscr{G}$ is a covering graph. The spectral properties of the Laplacian on general covering graphs are reported in [2]. In this paper, we focus on the covering graphs $G \in \mathscr{G}$ and investigate the spectrum of L_G in detail. Give an $n \in \mathbb{N}$ and r satisfying (1.1), we define the following subset of $V(G)$, which we call a *cell* of G:

$$V_{n,r}(G) = \{(n+m,s) \in V(G) \mid m = 0,1,\cdots,r-1, s = 0,1\}.$$

We note that the following relationships hold:

$$V(G) = \bigcup_{n\in\mathbb{Z}} V_{nr,r}(G), \quad V_{nr,r}(G) \cap V_{mr,r}(G) = \emptyset \quad (n \neq m).$$

We use $s(r)$ to denote the number of end vertices in the cell $V_{1,r}(G)$:

$$s(r) = \#\{(n,1) \in V(G) \mid n = 1,2,\cdots,r\}.$$

For instance, we have $r(G_{1,0}) = 1$, $s(G_{1,0}) = 0$; $r(G_{1,1}) = 1$, $s(G_{1,1}) = 1$ and $r(G_{2,1}) = 2$, $s(G_{2,1}) = 1$.

We say that G and H are isomorphic and write this as $G \simeq H$ if there exists a bijection $\phi : V(G) \longrightarrow V(H)$ such that $\phi(x) \sim \phi(y)$ if and only if $x \sim y$. For any graphs G, $H \in \mathscr{G}$, $G \simeq H$ holds if and only if there exists a number r satisfying (1.1) for both G and H such that $V_{1,r}(G) = V_{n,r}(H)$ for some $n \in \mathbb{Z}$. If $G \simeq H$, then $\sigma(L_G) = \sigma(L_H)$.

A graph $G \in \mathscr{G}$ is uniquely determined (up to isomorphism) by the number $s(r)$ of $(n,1) \in V_{1,r}(G)$ and the arrangement of $(n,1) \in V_{1,r}(G)$. In particular, the graph $G \in \mathscr{G}$ with $s(1) = 1$ has a single end vertex $(1,1) \in V(G)$ in the cell $V_{1,1}(G)$, and hence $G = G_{1,1}$. A graph $G \in \mathscr{G}$ with $s(2) = 1$ has a single end vertex $(n,1) \in V(G)$ in the cell $V_{1,2}(G)$; hence, G is isomorphic to $G_{2,1}$. Similarly, for any $r \in \mathbb{N}$ we can define a graph $G_{r,1} \in \mathscr{G}$ that has a single end vertex $(n,1) \in V(G)$ in the cell $V_{1,r}(G)$. Clearly, $G_{r,1}$ is uniquely determined up to isomorphism. For any $r \in \mathbb{N}$, we can define a graph $G_{r,r-1}$ such that $s(r) = r-1$, *i.e.* the number of end vertices $(n,1) \in V(G)$ in the cell $V_{1,r}(G)$ is equal to $r-1$.

On the other hand, in the case where $s(r) \neq 1$ or $s(r) \neq r-1$, there are a variety of graphs that are not isomorphic to each other. In general, for two graphs G and $\tilde{G}$ which are not isomorphic to each other, we cannot expect that $\sigma(L_G) = \sigma(L_{\tilde{G}})$. For instance, assuming $r = 4$ and $s(r) = 2$, we have two different graphs as follows: 1) if the end vertices $(n, 1) \in V_{1,4}(G_{4,2})$ are connected to alternate vertices, then the graph $G_{4,2} \in \mathscr{G}$ is isomorphic to $G_{2,1}$, and 2) the graph $\tilde{G}_{4,2} \in \mathscr{G}$ with end vertices $(1, 1)$ and $(2, 1) \in V_{1,4}(\tilde{G}_{4,2})$ satisfies $s(4) = 2$ but is not isomorphic to $G_{2,1}$. See Figures 3 and 4.



Figure 3: Graph $G_{4,2} \in \mathscr{G}$ has end vertices $(1, 1)$ and $(3, 1) \in V_{1,4}(G_{4,2})$.



Figure 4: Graph $\tilde{G}_{4,2} \in \mathscr{G}$ has end vertices $(1, 1)$ and $(2, 1) \in V_{1,4}(\tilde{G}_{4,2})$.

As we shall see later, the spectra of $L_{G_{2,1}}$ and $L_{G_{4,2}}$ are different. In particular, $L_{G_{2,1}}$ has zero as an eigenvalue whereas $L_{G_{4,2}}$ has no eigenvalue.

1.2 Results

Let $\mathscr{G}_\times = \mathscr{G} \setminus \{\mathbb{Z}\}$. We first identify spectral properties of L_G that hold for all graphs $G \in \mathscr{G}_\times$.

Theorem 1.1. Let $G \in \mathscr{G}_\times$. Then the following hold:

(1) $\sigma(L_G) \subset [-1, 1]$ is symmetric with respect to zero.

(2) There exists a constant $\epsilon > 0$ such that

$$(-\epsilon, 0) \cup (0, \epsilon) \subset \rho(L_G),$$

where $\rho(L_G)$ denotes the resolvent set of L_G.

(3) There is no eigenvalue in $\sigma(L_G) \setminus \{0\}$.

We will prove Theorem 1.1 in Section 3; here we will make two comments about the theorem: (i) The symmetry of $\sigma(L_G)$ with respect to zero in Theorem 1.1 (1) comes from the bipartiteness of $G \in \mathscr{G}$. If we remove condition (G_2) from the assumptions of Theorem 1.1, (1) will not hold true in general, as shown in Figure 5.



Figure 5: Let G be a graph satisfying (G_1) and (G_3) with $r = 2$ and suppose that $s(2) = 1$, $\deg(n, 0) = 3$, and $(1, 1) \in V_{1,2}(G)$ is connected to vertices $(1, 0)$ and $(2, 0)$. Then (G_2) does not hold. We observe that $1 \in \sigma(L_G)$, but $-1 \notin \sigma(L_G)$; hence, Theorem 1.1 (1) does not hold.

(ii) Theorem 1.1 (2) implies that for all graphs $G \in \mathscr{G}_\times$, L_G has a spectral gap around zero. The situation is different from the two-dimensional case. In Section 5, we will prove the following for graphs obtained from $\mathbb{Z}^2$ by adding pendant edges:

(a) there is an arrangement of pendant edges such that the Laplacian has no spectral gap (Theorem 5.1 (a));

(b) there is an arrangement of pendant edges such that the Laplacian has a spectral gap (Theorem 5.1 (b)).

In a companion paper [4], we study graphs obtained from the hexagonal lattice by adding pendant edges and establish results similar to that mentioned above. We believe that the spectra of the Laplacians on such graphs are related to the electronic structure of hydrogenated graphenes and graphane; see [1] for details.

Subsequently, we introduce a decomposition of the cells of a graph G in order to formulate the condition under which L_G has an eigenvalue. From Theorem 1.1 (3), we know that only the zero eigenvalue can exist: if it exists, then it is an isolated eigenvalue of L_G. For any r satisfying (1.1), we define

$$U_r(G) = \{(n, 0) \in V_{1,r}(G) \mid (n, 1) \notin V_{1,r}(G)\}.$$

Without loss of generality, we can assume that $(1, 1) \in V_{1,r}(G)$ because there exists a graph $\widetilde{G}$ isomorphic to G such that $(1, 1) \in V_{1,r}(\widetilde{G})$. Letting $\tilde{p}(r) = \#U_r(G)$, we can write

$$U_r(G) = \{(n_1, 0), (n_2, 0), \cdots, (n_{\tilde{p}(r)}, 0)\}$$

with $1 < n_1 < n_2 < \cdots < n_{\tilde{p}(r)}$. Then there exist a $p(r)$ with $1 \le p(r) \le \tilde{p}(r)$ and $l_1, l_2, \cdots, l_{p(r)} \ge 1$ such that $U_r(G)$ can be decomposed into the connected subgraphs U_i $(i = 1, 2, \cdots, p(r))$:

$$U_r(G) = \bigcup_{i=1}^{p(r)} U_i, \tag{1.2}$$

where

$$U_i = \left\{ \left(m_{1+\sum_{j=1}^{i-1} l_j}, 0 \right), \left(m_{2+\sum_{j=1}^{i-1} l_j}, 0 \right), \cdots, \left(m_{\sum_{j=1}^{i} l_j}, 0 \right) \right\} \tag{1.3}$$

with

$$m_{t+1} - m_t = 1, \quad t = 1 + \sum_{j=1}^{i-1} l_j, 2 + \sum_{j=1}^{i-1} l_j, \cdots, l_i - 1 + \sum_{j=1}^{i-1} l_j$$

$$m_{t+1} - m_t > 1, \quad t = \sum_{j=1}^{i} l_j, \ i = 1, 2, \cdots, p(r) - 1$$

and $\sum_{i=1}^{p(r)} l_i = \tilde{p}(r)$. See Figure 6 for an example. We use ●, ○ and △ to denote the vertices $(n, 0) \in U_r(G)$, $(n, 0) \notin U_r(G)$ and $(n, 1) \in V_{1,r}(G)$, respectively. Clearly, this decomposition is unique.



Figure 6: Let $G \in \mathscr{G}$ satisfy (1.1) with $r = 6$, and let $(1, 1), (5, 1) \in V_{1,6}(G)$ be the end vertices of G. Then the decomposition of $U_6(G)$ is $U_6(G) = U_1 \cup U_2$ with $U_1 = \{(2, 0), (3, 0), (4, 0)\}$ and $U_2 = \{(6, 0)\}$.

We have the following theorem:

Theorem 1.2. Let $G \in \mathscr{G}_\times$ and let $U_r(G) = \bigcup_{i=1}^{p(r)} U_i$ be the decomposition defined in (1.2). Then zero is an eigenvalue of L_G if and only if there exists an $i \in \{1, 2, \cdots, p(r)\}$ such that $l_i = \#U_i$ is odd.

We will prove Theorem 1.2 in Subsection 3.3.

Because, by virtue of (G$_3$), L_G is translation invariant for any $G \in \mathscr{G}$, we can produce more detailed information about the spectrum of L_G. For an r that satisfies (1.1), we define the operator $T_r : \ell^2(V(G)) \longrightarrow \ell^2(V(G))$ as

$$(T_r\psi)(n,s) = \psi(\tau_r(n,s)), \quad \psi \in \ell^2(V(G)),$$

where $\tau_r(n,s) = (n+r,s)$. We can show that L_G commutes with T_r, *i.e.* $[L_G, T_r] = 0$ because for any $\psi \in \ell^2(V(G))$,

$$\begin{aligned}(T_rL_G\psi)(n,s) = (L\psi)(\tau_r(n,s)) &= \frac{1}{\deg(\tau_r(n,s))}\sum_{y\sim\tau_r(n,s)}\psi(y)\\ &= \frac{1}{\deg(n,s)}\sum_{\tau_r^{-1}(y)\sim(n,s)}\psi(y) = \frac{1}{\deg(n,s)}\sum_{\tilde{y}\sim(n,s)}\psi(\tau_r(\tilde{y}))\\ &= (L_GT_r\psi)(n,s),\end{aligned}$$

where we have used the fact that $\deg(\tau_r(n,s)) = \deg(n,s)$ and set $\tilde{y} = \tau_r(y)$.

Let $X_r(G) = \{(n,0) \in V_{1,r}(G) \mid (n,0) \notin U_r(G)\}$ and set

$$X_r(G) = \{(i_1,0),(i_2,0),\cdots,(i_{s(r)},0)\}, \quad i_1 < i_2 < \cdots < i_{s(r)}. \tag{1.4}$$

As mentioned above, we can assume that $i_1 = 1$ without loss of generality. We define an identification $\iota : V(G) \longrightarrow \mathbb{Z}\times\{1,2,\cdots,r+s(r)\}$ with

$$\iota(nr+j,s) = \begin{cases}(n,j) & \text{if } j = 1,\cdots,r \text{ and } s = 0,\\ (n,r+p) & \text{if } j = i_p,\ p = 1,2,\cdots,s(r) \text{ and } s = 1.\end{cases}$$

for $(nr+j,s) \in V_{nr,r}(G)$, and we define a unitary operator $\mathscr{J} : \ell^2(V(G)) \longrightarrow \ell^2(\mathbb{Z};\mathbb{C}^{r+s(r)})$ as

$$(\mathscr{J}\psi)(n) = \begin{pmatrix}\psi_1(n)\\ \vdots\\ \psi_{r+s(r)}(n)\end{pmatrix}$$

with $\psi_i(n) = \psi(\iota^{-1}(n,i))$ $(i = 1,2,\cdots,r+s(r))$. The inner product of $\ell^2(\mathbb{Z};\mathbb{C}^{r+s(r)})$ is

$$\langle\mathscr{J}\psi|\mathscr{J}\phi\rangle := \sum_{n\in\mathbb{Z}}\langle\mathscr{J}\psi(n), D\mathscr{J}\phi(n)\rangle_{\mathbb{C}^{r+s(r)}}$$

with $D = \operatorname{diag}(d_1,d_2,\cdots,d_{r+s(r)})$ and

$$d_i = \deg(\iota^{-1}(n,i)), \quad i = 1,\cdots,r+s(r).$$

Note that d_i is independent of $n \in \mathbb{Z}$ because if $\iota^{-1}(n,i) = (nr+j,s)$, then $\iota^{-1}(n+1,i) = ((n+1)r+j,s) = \tau_r(nr+j,s)$. Let $\mathcal{F} : \ell^2(\mathbb{Z};\mathbb{C}^{r+s(r)}) \longrightarrow L^2([-\pi,\pi];\mathbb{C}^{r+s(r)})$ be the discrete Fourier transformation:

$$(\mathcal{F}\mathcal{J}\psi)(k) = \begin{pmatrix} \widehat{\psi}_1(k) \\ \vdots \\ \widehat{\psi}_{r+s(r)}(k) \end{pmatrix}, \quad k \in [-\pi,\pi],$$

where $\widehat{f}(k) = (2\pi)^{-1/2}\sum_{n\in\mathbb{Z}} e^{-ikn} f(n)$. We define the quasi momentum operator P as

$$P = (\mathcal{F}\mathcal{J})^{-1} M_k (\mathcal{F}\mathcal{J}),$$

where $M_{g(k)}$ denotes the multiplication operator by a function $g(k)$ on $L^2([-\pi,\pi];\mathbb{C}^{r+s(r)})$. Clearly, we have $\sigma(P) = [-\pi,\pi]$. We have the following lemma:

Lemma 1.3. Let r satisfy (1.1). Then $T_r = e^{iP}$.

Proof. Since we know from the functional calculus that $e^{iP} = (\mathcal{F}\mathcal{J})^{-1} M_{e^{ik}} (\mathcal{F}\mathcal{J})$, it suffices to prove that $(\mathcal{F}\mathcal{J})T_r = M_{e^{ik}}(\mathcal{F}\mathcal{J})$. Because

$$e^{ik}\widehat{\psi}_i(k) = (2\pi)^{-1/2}\sum_{n\in\mathbb{Z}} e^{-ik(n-1)}\psi_i(n) = \widehat{\psi_i(\cdot+1)}(k),$$

we have

$$(M_{e^{ik}}(\mathcal{F}\mathcal{J})\psi)(k) = \begin{pmatrix} \widehat{\psi_1(\cdot+1)}(k) \\ \vdots \\ \widehat{\psi_{r+s(r)}(\cdot+1)}(k) \end{pmatrix} = (\mathcal{F}\mathcal{J}T_r\psi)(k),$$

where we have used the following:

$$\begin{aligned} \psi_i(n+1) &= \psi(\iota^{-1}(n+1,i)) \\ &= \begin{cases} \psi((n+1)r+j,0), & i = j \ (j = 1,\cdots,r) \\ \psi((n+1)r+j,1), & i = r+p \ (j = i_p,\ p = 1,\cdots,s(r)) \end{cases} \end{aligned}$$

and

$$\psi((n+1)r+j,s) = \psi(\tau_r(nr+j,s)) = (T_r\psi)(nr+j,s) = (T_r\psi)_i(n).$$

□

From Lemma 1.3, we know that T_r is diagonalized by $\mathcal{U} := (\mathcal{F}\mathcal{J})^{-1}$ and $\sigma(T_r) = \{e^{ik} \mid k \in [-\pi,\pi]\}$. These facts imply that $\mathcal{U}$ can decompose L_G into a direct integral over the spectrum $[-\pi,\pi]$ of the quasi momentum operator P:

Proposition 1.4. Let $G \in \mathscr{G}$ and let r satisfy (1.1). Then we have

$$\mathcal{U}^{-1} L_G \mathcal{U} = \int_{[-\pi,\pi]}^{\oplus} L_k dk, \tag{1.5}$$

where L_k is an operator on $\mathcal{H}_k = \mathbb{C}^{r+s(r)}$ ($k \in [-\pi, \pi]$) and

$$\mathcal{U}^{-1} \ell^2(V(G)) = \int_{[-\pi,\pi]}^{\oplus} \mathcal{H}_k dk.$$

In Subsetion 2.1 we provide an explicit formula for L_k in Proposition 1.4. Let r satisfy (1.1) and $U_r(G) = \bigcup_{i=1}^{p(r)} U_i$ be the decomposition in (1.2). Then we use $q(r)$ to denote the number of the U_i for which $l_i = \# U_i$ is odd:

$$q(r) = \#\{U_i \mid l_i \text{ is odd}\}. \tag{1.6}$$

Proposition 1.5. Let r satisfy (1.1). Then

$$q(r) = \dim \ker L_k \tag{1.7}$$

for any $k \in [-\pi, \pi]$.

From Proposition 1.5, we know that the multiplicity of the zero eigenvalue of L_k is independent of $k \in [-\pi, \pi]$ and equal to $q(r)$. Thus, we know that there are at most $r + s(r) - q(r)$ non-zero eigenvalues of L_k. Let $\lambda_i(k)$ ($i = 1, 2, \cdots, r + s(r) - q(r)$) be the non-zero eigenvalues of L_k such that

$$\lambda_1(k) \geq \lambda_2(k) \geq \cdots \geq \lambda_{r+s(r)-q(r)}(k).$$

Note that

$$\sigma(L_G) \setminus \{0\} = \bigcup_{i=1}^{r+s(r)-q(r)} \{\lambda_i(k) \mid k \in [-\pi, \pi]\}. \tag{1.8}$$

Because the number of end vertices equals the number of vertices connected to end vertices, $q(r)$ is even (resp. odd) if $r + s(r)$ is even (resp. odd). Hence we know that $r + s(r) - q(r)$ is always even. Thus, we can define $t(r) \in \mathbb{N}$ as

$$t(r) = \frac{1}{2}(r + s(r) - q(r)). \tag{1.9}$$

We are now in a position to state our main result.

Theorem 1.6. Let $G \in \mathscr{G}_\times$ and $\lambda_i(k)$ $(i = 1, 2, \cdots, r + s(r) - q(r))$ be as given above. Then

$$\sigma(L_G) \setminus \{0\} = \bigcup_{i=1}^{t(r)} [a_i, b_i] \cup [-b_i, -a_i],$$

where

$$a_i = \begin{cases} \lambda_i(\pi) & \text{if } i \text{ is odd} \\ \lambda_i(0) & \text{if } i \text{ is even} \end{cases}, \quad b_i = \begin{cases} \lambda_i(0) & \text{if } i \text{ is odd} \\ \lambda_i(\pi) & \text{if } i \text{ is even} \end{cases}$$

for $i = 1, \cdots, t(r)$.

This theorem has the following corollary:

Corollary 1.7. Let $G \in \mathscr{G}_\times$. Then $\sigma(L_G)$ has at least one and at most $2t(r) - 1 (= r + s(r) - q(r))$ spectral gaps.

This corollary provides an estimate of the number of spectral gaps. It remains an open problem to determine the exact number of spectral gaps.

The rest of this paper is organized as follows. In Section 2, we present the proof of Proposition 1.4 with the explicit formula for L_k and derive the characteristic equation for L_k. In Section 3, we prove Proposition 1.5, parts (1) and (2) of Theorem 1.1 and Theorem 1.2. In Section 4, we prove Theorem 1.6 and part (3) of Theorem 1.1.

2 Characteristic equation

In subsequent sections, unless otherwise specified, we take it for granted that $G \in \mathscr{G}_\times$.

2.1 Decomposition of the transition operator

In this subsection, we prove Proposition 1.4. By definition, we know that

$$(\mathcal{J} L_G \psi)(n) = \begin{pmatrix} (L_G\psi)_1(n) \\ \vdots \\ (L_G\psi)_{r+s(r)}(n) \end{pmatrix} = \begin{pmatrix} \frac{1}{d_1}(\psi_r(n-1) + \psi_2(n) + \phi_1(n)) \\ \frac{1}{d_2}(\psi_1(n) + \psi_3(n) + \phi_2(n)) \\ \vdots \\ \frac{1}{d_{r-1}}(\psi_{r-2}(n) + \psi_r(n) + \phi_{r-1}(n)) \\ \frac{1}{d_r}(\psi_{r-1}(n) + \psi_1(n+1) + \phi_r(n)) \\ \psi_{i_1}(n) \\ \vdots \\ \psi_{i_{s(r)}}(n) \end{pmatrix},$$

where

$$\phi_i(n) = \begin{cases} \psi_{r+p}(n), & i = i_p \\ 0, & i \neq i_p \end{cases}, \quad p = 1, 2, \cdots, s(r).$$

Hence, we have

$$(\mathcal{F}\mathcal{J}L_G\psi)(k) = \begin{pmatrix} \frac{1}{d_1}(e^{-ik}\hat{\psi}_r(k) + \hat{\psi}_2(k) + \hat{\phi}_1(k)) \\ \frac{1}{d_2}(\hat{\psi}_1(k) + \hat{\psi}_3(k) + \hat{\phi}_2(k)) \\ \vdots \\ \frac{1}{d_{r-1}}(\hat{\psi}_{r-2}(k) + \hat{\psi}_r(k) + \hat{\phi}_{r-1}(k)) \\ \frac{1}{d_r}(\hat{\psi}_{r-1}(k) + e^{ik}\hat{\psi}_1(k) + \hat{\phi}_r(k)) \\ \hat{\psi}_{i_1}(k) \\ \vdots \\ \hat{\psi}_{i_{s(r)}}(k) \end{pmatrix} = L_k(\mathcal{F}\mathcal{J}\psi)(k),$$

where

$$L_k = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} \tag{2.1}$$

with

$$L_{11} = \begin{pmatrix} 0 & \frac{1}{d_1} & 0 & \cdots & 0 & \frac{e^{-ik}}{d_1} \\ \frac{1}{d_2} & 0 & \ddots & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \ddots & \ddots & \frac{1}{d_{r-1}} \\ \frac{e^{ik}}{d_r} & 0 & \cdots & 0 & \frac{1}{d_r} & 0 \end{pmatrix}, \tag{2.2}$$

$$L_{22} = \begin{pmatrix} 0 & \frac{1}{d_{r+1}} & 0 & \cdots & 0 \\ \frac{1}{d_{r+2}} & 0 & \ddots & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & \frac{1}{d_{r+s(r)-1}} \\ 0 & \cdots & 0 & \frac{1}{d_{r+s(r)}} & 0 \end{pmatrix}. \tag{2.3}$$

Here $L_{12} = (b_{i.j})$ and $L_{21} = (c_{i,j})$ are given by

$$b_{i,j} = \begin{cases} \frac{1}{d_i}, & i = i_p,\ j = p, \quad p = 1, \cdots, s(r) \\ 0, & \text{otherwise}, \end{cases} \tag{2.4}$$

$$c_{i,j} = \begin{cases} \frac{1}{d_{r+i}}, & i = p,\ j = i_p, \quad p = 1, \cdots, s(r) \\ 0, & \text{otherwise}. \end{cases} \tag{2.5}$$

Thus, we know that $\mathcal{U}^{-1}L_G\mathcal{U}\cdot(\mathcal{U}^{-1}\psi)(k)=L_k(\mathcal{U}^{-1}\psi)(k)$, which establishes Proposition 1.4.

We conclude this subsection with the following lemma:

Lemma 2.1. Let $k=0$. Then L_0 is a stochastic matrix, *i.e.* the elements of L_0 are all nonnegative and the sum of each row (resp. column) of L_0 is 1.

2.2 Characteristic equation

Let

$$\mathcal{U}^{-1}L_G\mathcal{U}=\int_{[-\pi,\pi]}^{\oplus}L_k dk$$

be the decomposition in (1.5). In this subsection, we investigate the characteristic polynomial for L_k and prove the following:

Proposition 2.2. Let $G\in\mathscr{G}$. Then there exist a positive constant $A_{(r+s(r)-\star)/2}$ and nonnegative constants A_m $(m=0,1,2,\ldots,(r+s(r)-\star)/2-1)$ independent of $k\in[-\pi,\pi]$ such that $\lambda=\lambda(k)$ is an eigenvalue of L_k if and only if

$$\sum_{m=0}^{(r+s(r)-\star)/2}A_m\lambda^{2m+\star}-2(\cos k)\lambda^{s(r)}=0, \tag{2.6}$$

where

$$\star=\begin{cases}0, & \text{if } r+s(r) \text{ is even,}\\ 1, & \text{if } r+s(r) \text{ is odd.}\end{cases} \tag{2.7}$$

In this proposition, $r+s(r)-\star$ is always even; hence, $(r+s(r)-\star)/2\in\mathbb{N}$. To prove this proposition, we introduce a directed weighted graph $G_k=(V(G_k),A(G_k))$ for L_k as follows. We use the ordered pair $[x,y]$ of two vertices x and y to denote the directed edge whose initial and terminal vertices are x and y, respectively. The set of vertices $V(G_k)$ of G_k is $V(G_k)=\{1,2,\cdots,r+s(r)\}$ and the set of directed edges of G_k is $A(G_k)=\{[i,i']\mid i\sim i',\ i,i'\in V(G_k)\}$, where $i\sim i'$ if i and i' satisfy one of the following three conditions:

(a) There exists an n such that $\iota^{-1}(n,i)\sim\iota^{-1}(n,i')$ as vertices of G.

(b) $(i,i')=(1,r)$ or $(i,i')=(r,1)$.

(c) $i=i'$, $i=1,2,\cdots,r+s(r)$.

Note that in the case of (c), the edge $[i, i]$ is a loop. We define the weight $a_{i,i'}$ of the edge $[i, i']$ as

$$a_{i,i'} = \begin{cases} -d_i\lambda & \text{if } i = i', \\ 1 & \text{if } \iota^{-1}(n, i) \sim \iota^{-1}(n, i'), \\ e^{ik} & \text{if } (i, i') = (r, 1), \\ e^{-ik} & \text{if } (i, i') = (1, r) \end{cases}$$

with $\lambda \in \mathbb{C}$. As a notational convention, we set $a_{i,i'} = 0$ if $[i, i'] \notin A(G_k)$. We define square matrices of order $r + s(r)$ by

$$A_k(\lambda) = A_k - \lambda D, \quad \lambda \in \mathbb{C},$$

where $A_k = (a_{i,i'})$ and $D = \mathrm{diag}(d_1, \cdots, d_{r+s(r)})$. We observe that from direct calculation,

$$A_k(\lambda) = (a_{i,i'}).$$

We now establish

Lemma 2.3. λ is an eigenvalue of L_k if and only if $A_k(\lambda)$ has a zero eigenvalue.

Proof. Since $DL_k = A_k$, we have

$$A_k(\lambda) = D(L_k - \lambda). \tag{2.8}$$

λ is an eigenvalue of L_k if and only if $\det(L_k - \lambda) = 0$. Because D is invertible and $\det(L_k - \lambda) = \det(D^{-1}) \cdot \det(A_k(\lambda))$, we have the desired result. $\square$

From Lemma 2.3, we know that the solution of the equation

$$\det(A_k(\lambda)) = 0$$

is an eigenvalue of L_k. Note that

$$\det(A_k(\lambda)) = \sum_{\sigma \in \mathfrak{S}_{r+s(r)}} A_\sigma, \tag{2.9}$$

where $\mathfrak{S}_n$ denotes the set of all permutations of degree n and

$$A_\sigma = \mathrm{sgn}(\sigma) \prod_{i=1}^{r+s(r)} a_{i,\sigma(i)}, \quad \sigma \in \mathfrak{S}_{r+s(r)}.$$

For $\sigma \in \mathfrak{S}_{r+s(r)}$, there exist integers $1 \le n \le r+s(r)$ and $n_j \ge 1$ $(j = 1, \cdots, n)$ such that $r + s(r) = \sum_{j=1}^{n} \mu_j$ and

$$\sigma = (i_1^{(1)}, \cdots, i_{\mu_1}^{(1)}) \cdots (i_1^{(n)}, \cdots, i_{\mu_n}^{(n)}), \tag{2.10}$$

where, for $\mu_j \ge 2$, $(i_1^{(j)}, \cdots, i_{\mu_j}^{(j)})$ denotes a cycle such that $i_1^{(j)}, \cdots, i_{\mu_j}^{(j)}$ satisfy

$$\sigma(i_1^{(j)}) = i_2^{(j)}, \quad \sigma(i_2^{(j)}) = i_3^{(j)}, \cdots, \sigma(i_{\mu_j}^{(j)}) = i_1^{(j)},$$

and for $\mu_j = 1$, $\sigma(i_1^{(j)}) = i_1^{(j)}$. For $\sigma \in \mathfrak{S}_{r+s(r)}$, we define n disjoint graphs $G_k^{(j)}$ $(j = 1, 2, \cdots, n)$ by

$$V(G_k^{(j)}) = \{i_1^{(j)}, \cdots, i_{\mu_j}^{(j)}\}, \quad A(G_k^{(j)}) = \{[i_\nu, \sigma(i_\nu)] \mid \nu = 1, \cdots, \mu_j\}. \tag{2.11}$$

We say that a finite directed graph $H = (V(H), E(H))$ with vertices $V(H) = \{h_1, \cdots, h_m\}$ is a cycle of length m if

$$A(H) = \{[h_1, h_2], [h_2, h_3], \cdots, [h_m, h_1]\}.$$

In particular, we call a cycle of length 1 a loop. We use $H_1 \dot\cup H_2$ to denote the disjoint union of two disjoint graphs H_1 and H_2, i.e. a graph whose vertices and edges are $V(H_1 \dot\cup H_2) = V(H_1) \cup V(H_2)$ and $A(H_1 \dot\cup H_2) = A(H_1) \cup A(H_2)$, respectively. We say that H is a spanning subgraph of G if $V(H) = V(G)$ and $A(H) \subset A(G)$. From definition, we know that $G_k^{(i)}$ $(j = 1, \cdots, n)$ are cycles, and if $\mu_j = 1$, then $G_k^{(j)}$ is a loop. Because $r + s(r) = \sum_{j=1}^{n} \mu_j$ and the graphs $G_k^{(j)}$ $(j = 1, 2, \cdots, n)$ are disjoint, we have $V(G_k) = \cup_{j=1}^{n} V(G_k^{(j)})$. Thus, we know that each $\sigma \in \mathfrak{S}_{r+s(r)}$ corresponds to a disjoint union $G_\sigma = G_k^{(1)} \dot\cup \cdots \dot\cup G_k^{(n)}$ with $G_k^{(j)}$ defined by (2.11). Let $\mathscr{G}_k$ be the set of graphs $\tilde{G}$ that satisfy $V(\tilde{G}) = V(G_k)$. Then we define a map $Q : \mathfrak{S}_{r+s(r)} \to \mathscr{G}_k$ as

$$Q(\sigma) = G_\sigma, \quad \sigma \in \mathfrak{S}_{r+s(r)}.$$

From its construction, the map Q is injective. Conversely, we suppose that disjoint graphs $G_k^{(j)}$ $(j = 1, 2, \cdots, n)$ are cycles with vertices $\{i_1^{(j)}, \cdots, i_{\mu_j}^{(j)}\} \subset V(G_k)$ such that $r + s(r) = \sum_{j=1}^{n} \mu_j$. Then the disjoint union $G_k^{(1)} \dot\cup \cdots \dot\cup G_k^{(n)}$ satisfies $V(G_k) = \cup_{j=1}^{n} V(G_k^{(j)})$. Let $\sigma \in \mathfrak{S}_{r+s(r)}$ be a permutation defined by (2.10). Then we have $Q(\sigma) = G_\sigma = G_k^{(1)} \dot\cup \cdots \dot\cup G_k^{(n)}$. Hence, we know that Q is bijective. The following lemma also holds:

Lemma 2.4. Let $\sigma \in \mathfrak{S}_{r+s(r)}$, A_σ and $G_\sigma = G_k^{(1)} \dot\cup \cdots \dot\cup G_k^{(n)}$ be as above. Then $A_\sigma \ne 0$ if and only if G_σ is a spanning subgraph of G_k.

Proof. It is clear that $A_\sigma \neq 0$ if and only if $a_{i,\sigma(i)} \neq 0$ for any $i \in \{1, 2, \cdots, r + s(r)\}$. Because, by definition, $a_{i,\sigma(i)} \neq 0$ if and only if $[i, \sigma(i)] \in A(G_k)$, we have the desired result. $\square$

We can now present the

Proof of Proposition 2.2. By virtue of Lemma 2.4, it suffices to consider the case where $G_\sigma = G_k^{(1)} \dot{\cup} \cdots \dot{\cup} G_k^{(n)}$ is a spanning subgraph of G_k.

Let H_k be a cycle given by

$$V(H_k) = \{1, 2, \cdots, r\}, \quad A(H_k) = \{[1, 2], [2, 3], \cdots, [r, 1]\}$$

We first consider a spanning subgraph $G_k^{(1)} \dot{\cup} \cdots \dot{\cup} G_k^{(n)}$ of G_k that includes a cycle H_k. Without loss of generality, we can set $H_k = G_k^{(1)}$. In this case, $G_k^{(j)}$ $(j \geq 2)$ cannot be cycles of length greater than 2 because $V(G_k^{(j)}) \subset \{r + 1, \cdots, r + s(r)\}$. Hence, we observe that $n = s(r) + 1$ and $G_k^{(J+1)}$ $(j = 1, \cdots, s(r))$ are loops with $V(G_k^{(J+1)}) = \{r + j\}$. Let $\sigma_+ \in \mathfrak{S}_{r+s(r)}$ be the permutation such that $G_{\sigma_+} = G_k^{(1)} \dot{\cup} \cdots \dot{\cup} G_k^{(s(r)+1)}$ with $G_k^{(j)}$ $(j = 1, \cdots, s(r) + 1)$ as mentioned above. Then we have $\sigma_+(1) = 2$, $\sigma_+(2) = 3$, $\cdots$, $\sigma_+(r) = 1$ and

$$\prod_{i=1}^{r} a_{i,\sigma_+(i)} = e^{ik}.$$

Because $\sigma_+(i) = i$ holds for $r + 1 \leq i \leq r + s(r)$, we obtain

$$\prod_{i=r+1}^{r+s(r)} a_{i,\sigma_+(i)} = (-\lambda)^{s(r)}.$$

Note that $\mathrm{sgn}\sigma_+ = (-1)^{r-1}$ since the length of the cycle H_k is r. Thus, we have

$$A_{\sigma_+} = (-1)^{r-1}(-\lambda)^{s(r)} e^{ik}.$$

Similarly, we set $\sigma_- \in \mathfrak{S}_{r+s(r)}$ such that $G_{\sigma_-} = G_k^{(1)} \dot{\cup} \cdots \dot{\cup} G_k^{(s(r)+1)}$ with $G_k^{(1)}$ replaced by H'_k, where H'_k is a cycle given by

$$V(H'_k) = \{1, 2, \cdots, r\}, \quad A(H'_k) = \{[1, r], [r, r - 1], \cdots, [2, 1]\}.$$

Then we have

$$A_{\sigma_-} = (-1)^{r-1}(-\lambda)^{s(r)} e^{-ik}.$$

Hence, we also have

$$A_{\sigma_+} + A_{\sigma_-} = 2(-1)^{r+s(r)-1}(\cos k)\lambda^{s(r)}. \tag{2.12}$$

Note that H_k (resp. H'_k) is the only cycle of length greater than 3 that contains $[1, r]$ (resp. $[r, 1]$); a cycle H_0 of length 2 that contains $[1, r]$ or $[r, 1]$ is given by

$$V(H_0) = \{1, r\}, \quad A(H) = \{[1, r], [r, 1]\}.$$

Hence, we know that if A_σ depends on $k \in [-\pi, \pi]$, then the edges $A(G_\sigma)$ of the spanning subgraph G_σ includes either H_k or H'_k. Let

$$\mathfrak{S}'_{r+s(r)} = \{\sigma \in \mathfrak{S}_{r+s(r)} \mid \sigma(1) = r,\, \sigma(r) = 1\}.$$

From the argument presented above, we have

$$\{\sigma \in \mathfrak{S}_{r+s(r)} \mid \sigma(1) = r \text{ or } \sigma(r) = 1\} = \mathfrak{S}'_{r+s(r)} \cup \{\sigma_+, \sigma_-\}.$$

Next we consider a cycle whose edges include both $[1, r]$ and $[r, 1]$. The cycle H_0 defined above is the only cycle that includes both $[1, r]$ and $[r, 1]$. We set $G_k^{(1)} = H_0$. Let $G_k^{(j)}$ $(j = 2, \cdots, n)$ be disjoint cycles and assume that $\tilde{G} := H_{(0)} \dot\cup G_k^{(2)} \dot\cup \cdots \dot\cup G_k^{(n)}$ is a spanning subgraph of G_k. Because there is no cycle of length grater than 3 that does not include edges $[1, r]$ and $[r, 1]$, the length of each $G_k^{(j)}$ $(j = 2, \cdots, n)$ is at most 2. We now claim that the number of loops included in the spanning subgraph $\tilde{G}$ is even if $r + s(r)$ is even. Note that if $r+s(r)$ is even, then $\#V(G_k)\backslash V(H_0) = r+s(r)-2$, which is even. If the spanning subgraph $\tilde{G}$ has $2\nu + 1$ loops, then cycles of length at most 2 must be constructed from the remaining $r + s(r) - 2 - (2m + 1)$ vertices. Because $r + s(r) - 2 - (2m + 1)$ is odd, these cycles include an odd number of loops. Thus, the claim is proven. Similarly, we can prove that the number of loops included in the spanning subgraph $\tilde{G}$ is odd if $r + s(r)$ is odd. Let $2m + \star$ $(0 \le m \le (r + s(r) - 2 - \star)/2)$ be the number of loops included in the spanning subgraph $\tilde{G}$, where $\star$ is defined as given in (2.7). Then the number of cycles of length 2 included in the spanning subgraph $\tilde{G}$ is $(r + s(r) - (2m + \star))/2$. Let $\sigma_{\tilde{G}}$ be the permutation which corresponds to $\tilde{G}$. Then $\sigma_{\tilde{G}} \in \mathfrak{S}_{r+s(r)}$ and we have $\mathrm{sgn}\sigma_{\tilde{G}} = (-1)^{1+(r+s(r)-(2m+\star))/2}$. Because $\sigma_{\tilde{G}}(1) = r$ and $\sigma_{\tilde{G}}(r) = 1$, it follows that $a_{1,\sigma_{\tilde{G}(1)}} a_{r,\sigma_{\tilde{G}(r)}} = e^{-ik} \cdot e^{ik} = 1$. Hence, we can write

$$\begin{aligned}\sum_{\sigma \in \mathfrak{S}'_{r+s(r)}} A_\sigma &= \sum_{m=0}^{(r+s(r)-2-\star)/2} B_m (-\lambda)^{2m+\star} (-1)^{(r+s(r)-(2m+\star))/2} \\ &= \sum_{m=0}^{(r+s(r)-2-\star)/2} B_m (-1)^{(r+s(r)+\star-2m)/2} \lambda^{2m+\star} \qquad (2.13)\end{aligned}$$

with constants $B_m \ge 0$ that are independent of $k \in [-\pi, \pi]$.

Let us calculate A_σ for $\sigma \in \mathfrak{S}_{r+s(r)}$ such that $\sigma \notin \mathfrak{S}'_{r+s(r)}$ and $\sigma \neq \sigma_\pm$. In this case, we have $\sigma(1) \neq r$ and $\sigma(r) \neq 1$. By reasoning similar to that mentioned above, we can assume that there are $2m + \star$ loops and $(r + s(r) - (2m + \star))/2$ cycles of length 2 in the spanning subgraph that corresponds to σ, where $0 \leq m \leq (r + s(r) - \star)/2$. Hence, we obtain

$$\begin{aligned}\sum_{\sigma:\sigma(1)\neq r,\sigma(r)\neq 1} A_\sigma &= \sum_{m=0}^{(r+s(r)-\star)/2} C_m(-\lambda)^{2m+\star}(-1)^{(r+s(r)-(2m+\star))/2} \\ &= \sum_{m=0}^{(r+s(r)-\star)/2} C_m(-1)^{(r+s(r)+\star-2m)/2}\lambda^{2m+\star} \qquad (2.14)\end{aligned}$$

with $C_m \geq 0$ independent of $k \in [-\pi, \pi]$. We observe that

$$C_{(r+s(r)-\star)/2} = \prod_{i=1}^{r+s(r)} d_i > 0 \qquad (2.15)$$

because the cycles included in the corresponding spanning subgraph are loops.

From (2.12), (2.13) and (2.14), we have

$$\begin{aligned}\sum_{\sigma\in\mathfrak{S}_{r+s(r)}} A_\sigma &= C_{(r+s(r)-\star)/2}(-1)^\star\lambda^{r+s(r)} \\ &+ \sum_{m=0}^{(r+s(r)-2-\star)/2} (B_m + C_m)(-1)^{(r+s(r)+\star-2m)/2}\lambda^{2m+\star} \\ &+ 2(-1)^{r+s(r)-1}(\cos k)\lambda^{s(r)}\end{aligned}$$

Based on Lemma 2.3 and (2.9), we know that λ is an eigenvalue of L_k if and only if

$$\begin{aligned}&C_{(r+s(r)-\star)/2}\lambda^{r+s(r)} \\ &+ \sum_{m=0}^{(r+s(r)-2-\star)/2} (-1)^{-\star+(r+s(r)+\star-2m)/2}(B_m + C_m)\lambda^{2m+\star} \\ &- 2(-1)^{-\star+r+s(r)}(\cos k)\lambda^{s(r)} = 0.\end{aligned}$$

Let $A_{(r+s(r)-\star)/2} = C_{(r+s(r)-\star)/2}$ and

$$A_m = (-1)^{(r+s(r)-2m-\star)/2}(B_m + C_m), \quad 0 \leq m \leq (r + s(r) - 2 - \star)/2. \quad (2.16)$$

From (2.15), we know that $A_{(r+s(r)-\star)/2}$ is strictly positive. Since $-\star+r+s(r)$ is always even and $(-1)^{-\star+r+s(r)} = 1$, Proposition 2.2 is proven. □

2.3 Lowest degree of the characteristic polynomial

In this subsection, we prove the following proposition:

Proposition 2.5. Let A_m be as in Proposition 2.2 and let $q(r)$ be defined by (1.6). Then the following holds:

$$q(r) = \inf\{2m + \star \mid A_m \neq 0\}.$$

To prove this proposition, let $U_r(G) = \bigcup_{i=1}^{p(r)} U_i$, where

$$U_i = \left\{ \left(m_{1+\sum_{j=1}^{i-1} l_j}, 0 \right), \left(m_{2+\sum_{j=1}^{i-1} l_j}, 0 \right), \cdots, \left(m_{\sum_{j=1}^{i} l_j}, 0 \right) \right\}$$

is the decomposition defined in (1.2) and (1.3). Then we define a corresponding decomposition for G_k by

$$U_k = \bigcup_{i=1}^{p(r)} U_k^{(i)},$$

where

$$U_k^{(i)} = \left\{ m_{1+\sum_{j=1}^{i-1} l_j}, m_{2+\sum_{j=1}^{i-1} l_j}, \cdots, m_{\sum_{j=1}^{i} l_j} \right\}.$$

For each $U_k^{(i)}$, we define cycles $K_\alpha^{(i)}$ whose vertices are included in $U_k^{(i)}$ as follows: If $\# U_k^{(i)} = \# U_i$ is even, then we set

$$V(K_\alpha^{(i)}) = \{ m_{2\alpha-1+\sum_{j=1}^{i-1} l_j}, m_{2\alpha+\sum_{j=1}^{i-1} l_j} \}, \quad 1 \leq \alpha \leq m_{\sum_{j=1}^{i} l_j}/2.$$

If $\# U_k^{(i)} = \# U_i$ is odd, then we define a loop $K_1^{(i)}$ with $V(K_1^{(i)}) = \{ m_{1+\sum_{j=1}^{i-1} l_j} \}$ and cycles $K_\alpha^{(i)}$ of length 2 with

$$V(K_\alpha^{(i)}) = \{ m_{2\alpha+\sum_{j=1}^{i-1} l_j}, m_{1+2\alpha+\sum_{j=1}^{i-1} l_j} \}, \quad 1 \leq \alpha \leq (m_{\sum_{j=1}^{i} l_j} - 1)/2.$$

Let $1 = i_1 < i_2 < \cdots < i_{s(r)}$ be as in (1.4). Then, by definition, we know that $i_j \notin U_k$ and

$$[i_j, r+j], [r+j, i_j] \in A(G_k), \quad j = 1, \cdots, s(r).$$

We define cycles $H_k^{(j)}$ $(j = 1, \cdots, s(r))$ of length 2 with

$$V(H_k^{(j)}) = \{ i_j, r+j \}, \quad A(H_k^{(j)}) = \{ [i_j, r+j], [r+j, i_j] \}.$$

Since $V(G_k) \setminus (\cup_j V(H_k^{(j)})) = U_k$, we have a spanning graph

$$K = \left(\dot{\bigcup_j} H_k^{(j)} \right) \dot{\cup} \left(\dot{\bigcup_{i,\alpha}} K_\alpha^{(i)} \right). \tag{2.17}$$

By definition, K has exactly $q(r)$ loops. In general, K is not a unique spanning subgraph of G_k that has exactly $q(r)$ loops. Indeed, if $q(r) \geq 1$, then we can construct such a spanning subgraph of G_k as follows: Since $q(r) \geq 1$, there is a number i_0 such that $\# U_k^{(i)}$ is odd. Hence, by definition, $K_1^{(i_0)}$ is a loop with the vertex $m_{1+\sum_{j=1}^{i-1} l_j}$. Then there is a cycle $H_k^{(j_0)}$ with vertices $i_{j_0}, r + j_0$ such that

$$[i_{j_0}, m_{1+\sum_{j=1}^{i-1} l_j}] \in A(G_k).$$

Hence, we can define a spanning subgraph K' with exactly $q(r)$ loops by

$$K' = \left(\dot{\bigcup_{j \neq j_0}} H_k^{(j)} \right) \dot{\cup} \left(\dot{\bigcup_{(i,\alpha) \neq (i_0,1)}} K_\alpha^{(i)} \right) \dot{\cup} M,$$

where M is the disjoint union of a loop with vertex $r + j_0$ and a cycle with vertices i_{j_0} and $m_{1+\sum_{j=1}^{i-1} l_j}$.

Let

$$m_0 = \frac{1}{2}(q(r) - \star).$$

Since $q(r) - \star$ is always even, we know that m_0 is a nonnegative integer. In particular, $m_0 = 0$ if $q(r) = 0$ or $q(r) = 1$. We shall now prove

Lemma 2.6. The following hold:

(a) If $q(r) = 0$, then $|A_0| = 1$.

(b) If $q(r) \geq 1$, then $|A_{m_0}| > 2$.

In particular, we have

$$\inf\{2m + \star \mid A_m \neq 0\} \leq q(r). \tag{2.18}$$

Proof. Let a_σ denote the absolute value of the coefficient of A_σ. Then for any spanning subgraph $\tilde{K}$ of G_k which has exactly $q(r)$ loops, we have $A_{\sigma_{\tilde{K}}} = (\mathrm{sgn}\sigma_{\tilde{K}}) a_{\sigma_{\tilde{K}}} \lambda^{q(r)}$. We assume that $\sigma_{\tilde{K}} \neq \sigma_\pm$ if $q(r) = s(r)$. Because every $\tilde{K}$ has the same number of cycles of length 2 as K, we know that $\sigma_{\tilde{K}} = \sigma_K$. Hence we have $|A_{m_0}| = \sum_{\tilde{K}} a_{\sigma_{\tilde{K}}}$.

To prove (a), we assume that $q(r) = 0$. Then $\#U_k^{(i)}$ is even for $i = 1, \cdots, p(r)$ and all disjoint subgraphs included in K are cycles of length 2. Since all the spanning subgraphs of G_k that do not have loops include $H_k^{(j)}$ $(j = 1, \cdots, s(r))$, K is a unique spanning subgraph of G_k without loops. Hence $|A_0| = a_{\sigma_K} = 1$.

To prove (b), we assume that $q(r) \geq 1$. Then, as mentioned above, K and K' have exactly $q(r)$ loops. We know $a_{\sigma_K} \geq 2$, because $K_1^{(i_0)}$ is the loop with the vertex $m_{1+\sum_{j=1}^{i-1} l_j}$ and $\deg m_{1+\sum_{j=1}^{i-1} l_j} = 2$. Hence, $|A_{m_0}| \geq a_{\sigma_K} + a_{\sigma_{K'}} > 2$.

Because $q(r) = 2m_0 + \star$, (2.18) is obtained from (a) and (b). □

We use $\mathcal{K}(\{M^{(i)}\}_{i=1}^{s})$ to denote the set of all spanning subgraphs of G_k that include disjoint graphs $M^{(i)}$ $(i = 1, \cdots, s)$. Let $q(\tilde{G})$ be the number of loops included in a spanning subgraph $\tilde{G}$ of G_k. For any spanning subgraph $\tilde{G}$ in $\mathcal{K}(\{H_k^{(j)}\}_{j=1}^{s(r)})$, the lengths of the cycles included in $\tilde{G}$ are all less than or equal to 2 because $[1, r+1] \in A(H_k^{(1)})$. Hence, $q(\tilde{G})$ equals the number of loops with a vertex in U_k. If $l_i = \#U_k^{(i)}$ is odd, then $\tilde{G} \in \mathcal{K}(\{H_k^{(j)}\}_{j=1}^{s(r)})$ has at least one loop with a vertex of $U_k^{(i)}$; if $l_i = \#U_k^{(i)}$ is odd, then $\tilde{G}$ contains no loop with a vertex of $U_k^{(i)}$. Therefore, $q(r)$ is the minimum of the number of loops included in a spanning subgraph in $\mathcal{K}(\{H_k^{(j)}\}_{j=1}^{s(r)})$:

$$q(r) = \min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{H_k^{(j)}\}_{j=1}^{s(r)})\}. \tag{2.19}$$

Let $1 \leq j_1 < j_2 < \cdots < j_s \leq s(r)$ $(1 \leq s \leq s(r))$ and $i_0 = 1, \cdots, s$. Because $\mathcal{K}(\{H_k^{(j_i)}\}_{i=1}^{s}) \subset \mathcal{K}(\{H_k^{(j_i)}\}_{i \neq i_0})$, we have

$$\min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{H_k^{(j_i)}\}_{i \neq i_0})\} \leq \min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{H_k^{(j_i)}\}_{i=1}^{s})\}. \tag{2.20}$$

Indeed, the converse inequality of (2.20) holds as well as the following:

Lemma 2.7. Let $1 \leq j_1 < j_2 < \cdots < j_s \leq s(r)$ $(1 \leq s \leq s(r))$. Then

$$\min_{i_0=1,2,\cdots,s} \min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{H_k^{(j_i)}\}_{i \neq i_0})\} = \min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{H_k^{(j_i)}\}_{i=1}^{s})\}.$$

Before proving Lemma 2.7, we complete the proof of Proposition 2.5:

Proof. We only require to prove the converse of the inequality (2.18). From Lemma 2.7 and (2.19), we know that

$$\min_{s=1,\cdots,s(r)} \min_{1 \leq j_1 < \cdots < j_s \leq s(r)} \min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{H_k^{(j_i)}\}_{i=1}^{s})\} = q(r).$$

Note that the set of all spanning subgraphs of G_k is

$$\{G_{\sigma_+}, G_{\sigma_-}\} \cup \left[\bigcup_{s=1}^{s(r)} \bigcup_{1 \leq j_1 < \cdots < j_s \leq s(r)} \mathcal{K}(\{H_k^{(j_i)}\}_{i=1}^{s}) \right].$$

Since the number of loops included in a spanning subgraph $\tilde{G}$ equals the degree of $A_{Q^{-1}(\tilde{G})}$ in λ, we have

$$\inf\{2m + \star \mid A_m \neq 0\} \geq q(r).$$

This completes the proof of Proposition 2.5. □

Proof of Lemma 2.7. We prove the lemma for the case where $s = s(r)$; the other case can be similarly proven. It suffices to prove that

$$\min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{H_k^{(j)}\}_{j \neq j_0})\} \geq q(r). \tag{2.21}$$

If $\tilde{G} \in \mathcal{K}(\{H_k^{(j)}\}_{j \neq j_0}) \backslash \mathcal{K}(\{H_k^{(j)}\}_{j=1}^{s(r)})$, then $\tilde{G}$ includes the loop with the vertex $r + j_0$, because $\tilde{G}$ does not include $H_k^{(j_0)}$. Hence, we have

$$\mathcal{K}(\{H_k^{(j)}\}_{j \neq j_0}) = \mathcal{K}(\{H_k^{(j)}\}_{j=1}^{s(r)}) \cup \mathcal{K}(\{\tilde{H}_k^{(j)}\}_{j=1}^{s(r)}),$$

where $\tilde{H}_k^{(j)} = H_k^{(j)}$ $(j \neq j_0)$ and $\tilde{H}_k^{(j_0)}$ denotes the loop with vertex $r + j_0$. Thus, we observe that

$$\min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{H_k^{(j)}\}_{j \neq j_0})\} = \min\{q(r), q_0(r)\},$$

where

$$q_0(r) = \min\{q(\tilde{G}) \mid \tilde{G} \in \mathcal{K}(\{\tilde{H}_k^{(j)}\}_{j=1}^{s(r)})\}.$$

Therefore, (2.21) will be proven if the following holds:

$$q_0(r) \geq q(r). \tag{2.22}$$

We will only prove (2.22) for the case where $1 < j_0 < s(r)$; (2.22) can be proven in a similar manner for $i_0 = 1, s(r)$. The following cases are possible for $\tilde{G} \in \mathcal{K}(\{\tilde{H}_k^{(j)}\}_{j=1}^{s(r)})$:

(i) There exists an i_1 such that $i_{j_0} - 1 \in U_k^{(i_1)}$ and $i_{j_0} + 1 \in U_k^{(i_1+1)}$;

(ii) There exists an i_1 such that $i_{j_0} - 1 \in H_k^{(j_0-1)}$ and $i_{j_0} + 1 \in H_k^{(j_0+1)}$;

(iii) There exists an i_1 such that $i_{j_0} - 1 \in U_k^{(i_1)}$ and $i_{j_0} + 1 \in H_k^{(j_0+1)}$;

(iv) There exists an i_1 such that $i_{j_0}-1 \in H_k^{(j_0-1)}$ and $i_{j_0}+1 \in U_k^{(i_1)}$.

We consider case (i) first. Since there is no cycle of length grater than 2, we need only consider the following three subcases: $\tilde{G} \in \mathcal{K}(\{\tilde{H}_k^{(j)}\}_{j=1}^{s(r)})$ includes (a) the loop with the vertex $i_{j_0} \in V(\tilde{G})$, (b) the cycle with the vertices i_{j_0} and $i_{j_0}-1$, and (c) the cycle with the vertices i_{j_0} and $i_{j_0}+1$. If (a) holds, then so does the following:

$$q(\tilde{G}) \geq 2 + q(r). \tag{2.23}$$

Next, we consider the cases (b) and (c). We define

$$\tilde{U}_k^{(i)} := \begin{cases} U_k^{(i)}, & 1 \leq i \leq i_1 - 1, \\ U_k^{i_1} \cup \{i_{j_0}\} \cup U_k^{i_1+1}, & i = i_1, \\ U_k^{(i+1)}, & i \geq i_1 + 1. \end{cases}$$

Then we have

$$U_k \cup \{i_{j_0}\} = \bigcup_{i=1}^{p(r)-1} \tilde{U}_k^{(i)}.$$

Let

$$\tilde{q} = \#\{\tilde{U}_k^{(i)} \mid \#\tilde{U}_k^{(i)} \text{ is odd}\}.$$

By an argument similar to the above, we can show that

$$q(\tilde{G}) \geq \tilde{q} + 1. \tag{2.24}$$

Let us calculate $\tilde{q}$. Since $\#\tilde{U}_k^{(i_1)}$ is odd if and only if both $\#U_k^{(i_1)}$ and $\#U_k^{(i_1+1)}$ are even or both are odd, we have

$$\tilde{q} = \begin{cases} q(r) - 1, & \text{if both } \#U_k^{(i_1)} \text{ and } \#U_k^{(i_1+1)} \text{ are odd}, \\ q(r) + 1, & \text{if both } \#U_k^{(i_1)} \text{ and } \#U_k^{(i_1+1)} \text{ are even}, \\ q(r), & \text{otherwise}. \end{cases}$$

From (2.24), we know that if (b) or (c) holds, then

$$q(\tilde{G}) \geq q(r). \tag{2.25}$$

From (2.23) and (2.25), we obtain (2.22) for case (i).

For case (ii), we observe that $\tilde{G} \in \mathcal{K}(\{\tilde{H}_k^{(j)}\}_{j=1}^{s(r)})$ includes the loop with the vertex i_{j_0}. By an argument similar to the above, we have

$$q(r) + 2 = q_0(r).$$

This establishes (2.22) for case (ii).

By an argument similar to the above, we know that in case (iii) $\tilde{G} \in \mathcal{K}(\{\tilde{H}_k^{(j)}\}_{j=1}^{s(r)})$ must include include the cycle with vertices i_{j_0} and $i_{j_0}-1$. We set

$$\hat{U}_k^{(i)} := \begin{cases} U_k^{(i)}, & i \neq i_1, \\ U_k^{(i_1)} \cup \{i_{j_0}\}, & i = i_1 \end{cases}$$

and $\hat{q} = \#\{\hat{U}_k^{(i)} \mid \#\hat{U}_k^{(i)} \text{ is odd }\}$. Then $q(\tilde{G}) \geq \hat{q}+1$ holds. Since $\hat{U}_k^{(i)}$ is odd if and only if $U_k^{(i_1)}$ is even, we have

$$\hat{q} = \begin{cases} q(r)+1, & \text{if } U_k^{(i_1)} \text{ is even,} \\ q(r)-1, & \text{if } U_k^{(i_1)} \text{ is odd.} \end{cases}$$

This establishes (2.22) for case (iii), and (2.22) is proven for case (iv) holds in a similar manner. □

3 Spectrum

In this section, we prove parts (1) and (2) of Theorem 1.1, along with Theorem 1.2.

3.1 Symmetry of the spectrum

We first prove part (1) of Theorem 1.1. Let $\lambda \in \sigma(L_G) \setminus \{0\}$. It suffices to show that $-\lambda \in \sigma(L_G)$. From Proposition 2.2, $\lambda \in \sigma(L_G)$ if and only if there exists a $k \in [-\pi, \pi]$ such that (2.2) holds. Hence, we have

$$\sum_{m=0}^{(r+s(r)-\star)/2} A_m \lambda^{2m+\star} = 2(\cos k)\lambda^{s(r)}$$

for some $k \in [-\pi, \pi]$. Then

$$\begin{aligned} \sum_{m=0}^{(r+s(r)-\star)/2} A_m(-\lambda)^{2m+\star} &= (-1)^{\star} \sum_{m=0}^{(r+s(r)-\star)/2} A_m \lambda^{2m+\star} \\ &= 2(-1)^{s(r)+\star}(\cos k)(-\lambda)^{s(r)}. \end{aligned} \tag{3.1}$$

If $s(r)+\star$ is even, then (3.1) means that $-\lambda \in \sigma(L)$. If $s(r)+\star$ is odd, then

$$\sum_{m=0}^{(r+s(r)-\star)/2} A_m(-\lambda)^{2m+\star} = 2\cos(k \pm \pi)(-\lambda)^{s(r)},$$

which also implies that $-\lambda \in \sigma(L_G)$.

3.2 Spectral gap

Next, we prove part (2) of Theorem 1.1. From part (1) of Theorem 1.1, we know that $\rho(L_G)$ is symmetric with respect to 0. It suffices to show that

$$(0,\epsilon) \subset \rho(L_G) \tag{3.2}$$

for some (possibly small) $\epsilon > 0$. From Propositions 2.2 and 2.5, we know that $\lambda \in \rho(L_G)$ if and only if

$$\sum_{m=m_0}^{(r+s(r)-\star)/2} A_m \lambda^{2m+\star} \neq 2(\cos k)\lambda^{s(r)} \quad \text{for any } k \in [-\pi,\pi], \tag{3.3}$$

where $m_0 = (q(r)-\star)/2$. Suppose that $\lambda > 0$. Then (3.3) is equivalent to

$$\left(\sum_{m=m_0}^{(r+s(r)-\star)/2} \frac{A_m}{2}\lambda^{2m+\star-s(r)}\right)^2 > 1. \tag{3.4}$$

We first suppose that $q(r) = 0$. Then, from (a) of Lemma 2.6, we know that $\star = 0$ and the left hand side of (3.4) is

$$\left(\sum_{m=0}^{(r+s(r)-\star)/2} \frac{A_m}{2}\lambda^{2m-s(r)}\right)^2 = C_0\lambda^{-2s(r)} + O(\lambda^{4-2s(r)})$$

with $C_0 = (A_{m_0}/2)^2 = 1/4$. Since $s(r) \geq 1$, this establishes (3.4) for sufficiently small $\lambda > 0$.

Let us next assume that $q(r) \geq 1$. Then, from part (b) of Lemma 2.6, it follows that the left hand side of (3.4) is

$$\left(\sum_{m=m_0+1}^{(r+s(r)-\star)/2} \frac{A_m}{2}\lambda^{2m+\star-s(r)}\right)^2 = C_1\lambda^{2q(r)-2s(r)} + O(\lambda^{4+2q(r)-2s(r)}) \tag{3.5}$$

with $C_1 = (A_{m_0}/2)^2 > 1$. Since, by definition, $q(r) \leq p(r)$ and $p(r) \leq s(r)$, we know that $q(r) \leq s(r)$. If $q(r) = s(r)$, then the left hand side of (3.5) is equal to $C_1 + O(\lambda^4)$. In this case, since $C_1 > 1$, (3.4) holds for sufficiently small $\lambda > 0$. If $q(r) < s(r)$, then, because $2q(r) - 2s(r) \leq -2$ and $C_1 > 0$, (3.4) holds for sufficiently small $\lambda > 0$. Thus, (3.2) is proven.

3.3 Existence of the zero eigenvalue

In this subsection, we prove Theorem 1.2 and Proposition 1.5. From Propositions 2.2 and 2.5, we know that λ is an eigenvalue of L_k if and only if

$$\lambda^{q(r)} F(k, \lambda) = 0, \tag{3.6}$$

where $m_0 = (q(r) - \star)/2$ and

$$\begin{aligned} F(k, \lambda) &= \sum_{m=m_0}^{(r+s(r)-\star)/2} A_m \lambda^{2(m-m_0)} - 2(\cos k)\lambda^{s(r)-q(r)} \\ &= \sum_{m=0}^{(r+s(r)-q(r))/2} A_{m+m_0} \lambda^{2m} - 2(\cos k)\lambda^{s(r)-q(r)}. \end{aligned} \tag{3.7}$$

From Lemma 2.6, we have

$$F(k, 0) = A_{m_0} - 2\delta_{s(r),q(r)} \cos k \neq 0, \qquad k \in [-\pi, \pi],$$

where

$$\delta_{s(r),q(r)} = \begin{cases} 1, & \text{if } s(r) = q(r), \\ 0, & \text{if } s(r) > q(r). \end{cases}$$

From [5, Theorem XIII.85], we know that L_G has 0 as an eigenvalue if and only if the Lebesgue measure of the set

$$\{k \in [-\pi, \pi] \mid L_k \text{ has 0 as an eigenvalue}\}$$

is positive. Hence, we observe that $0 \in \sigma_{\mathrm{p}}(L_G)$ is equivalent to $q(r) \geq 1$. Since $q(r) \geq 1$ if and only if $\#U_i$ is odd for some $i \in \{1, 2, \cdots, p(r)\}$, Theorem 1.2 is proven.

To prove (1.7) in Proposition 1.5, we review some basic facts. Let L be a self-adjoint operator on a Hilbert space $\mathcal{L}$ and λ_0 an eigenvalue of L. Then the geometric multiplicity $m_{\mathrm{g}}(\lambda_0) = \dim \ker(L - \lambda_0)$ of λ_0 is equal to the algebraic multiplicity $m_{\mathrm{a}}(\lambda_0)$. If $\mathcal{L}$ is a finite dimensional space and λ_i are eigenvalues of L, then

$$\det(L - \lambda) = \prod_i (\lambda_i - \lambda)^{m_{\mathrm{a}}(\lambda_i)}$$

(see [3, p.42]). From this general fact, we know that (1.7) holds.

4 Band structure of the spectrum

4.1 Band functions

We consider the solution $\lambda = \lambda(k)$ of

$$F(k, \lambda) = 0, \tag{4.1}$$

where $F(k, \lambda)$ is a polynomial of degree $r + s(r) - q(r)$ as defined in (3.7). Let $\lambda_i(k)$ $(i = 1, \cdots, r + s(r) - q(r))$ be solutions of (4.1) and assume that

$$\lambda_1(k) \geq \lambda_2(k) \geq \cdots \geq \lambda_{r+s(r)-q(r)}.$$

Then

$$F(k, \lambda) = A_{(r+s(r)-q(r))/2} \prod_{i=1}^{r+s(r)-q(r)} (\lambda_i(k) - \lambda). \tag{4.2}$$

We also have the following lemmas:

Lemma 4.1. $\lambda_1(0) = 1$ and

$$\dim \ker(L_0 - \lambda_1(0)) = 1.$$

Proof. From Lemma 2.1, L_0 is a stochastic matrix. The well-known fact that the largest eigenvalue of a stochastic matrix is unique and equal to 1 yields the desired result. $\square$

Lemma 4.2. Let $\lambda_i(k)$ $(i = 1, \cdots, r + s(r) - q(r))$ be as described above. Then the following hold for $i = 1, 2, \cdots, r + s(r) - q(r)$:

(1) For any $k \in (-\pi, 0) \cup (0, \pi)$, $\dim \ker(L_k - \lambda_i(k)) = 1$. In particular, we have

$$\lambda_1(k) > \lambda_2(k) > \cdots > \lambda_{r+s(r)-q(r)}(k), \quad k \in (-\pi, 0) \cup (0, \pi). \tag{4.3}$$

(2) For $k = 0, \pm\pi$, $\dim \ker(L_k - \lambda_i(k)) \leq 2$.

We postpone the proof of Lemma 4.2 until the next subsection.

Because we know, from (3.7), that $F(k, \lambda) = F(-k, \lambda)$, we have

$$\lambda_i(-k) = \lambda_i(k), \quad k \in [-\pi, \pi]. \tag{4.4}$$

From (1.8) and (4.4), we have

$$\sigma(L_G) \setminus \{0\} = \bigcup_{i=1}^{r+s(r)-q(r)} \{\lambda_i(k) \mid k \in [0, \pi]\}. \tag{4.5}$$

Therefore, we only require to study the behaviour of the solutions $\lambda_i(k)$ for $k \in [0, \pi]$.

Lemma 4.3. Let $\lambda_i(k)$ $(i = 1, \cdots, r + s(r) - q(r))$ be as described above. Then λ_i is analytic in a region that includes $(0, \pi)$ and is continuous in $[0, \pi]$.

Proof. We first prove the first half of the lemma. From (4.2) and (4.3), we have

$$\frac{\partial}{\partial \lambda} F(k, \lambda)|_{\lambda=\lambda_i(k)} = -A_{(r+s(r)-q(r))/2} \prod_{j \neq i} (\lambda_j(k) - \lambda_i(k)) \neq 0, \quad k \in (0, \pi)$$

for $i = 1, \cdots, r + s(r) - q(r)$. Because $F(k, \lambda_i(k)) = 0$, we know from the implicit function theorem that $\lambda_i(k)$ is a unique analytic function in a region which includes $(0, \pi)$.

We next prove the second half of the lemma. From Lemma 4.2, $\lambda_i(0)$ is a discrete eigenvalue of multiplicity at most 2. Because $\{L_k\}_k$ is an analytic family in the sense of Kato for k near 0 (see [5, Theorem XII.13]), we know that $\lambda_i(k)$ is continuous at $k = 0$. The proof of the continuity at $k = \pi$ is similar. ☐

Now let

$$D(\lambda) = \frac{1}{2} \sum_{m=0}^{t(r)} A_{m+m_0} \lambda^{2m-s(r)+q(r)},$$

where $t(r)$ is defined in (1.9). We observe that $D(\lambda)$ is an even (resp. odd) function if r is odd (resp. even), because $s(r) - q(r)$ is even (resp. odd) if r is odd (resp. even). From (3.7) and (4.1), we know that $\lambda = \lambda(k)$ is a solution of (4.1) if and only if

$$D(\lambda(k)) = \cos k. \tag{4.6}$$

Lemma 4.4. Let $\lambda_i(k)$ $(i = 1, \cdots, r + s(r) - q(r))$ be as described above. Then:

(1) If r is even,

$$\lambda_{2t(r)+1-i}(k) = -\lambda_i(k), \quad k \in [0, \pi], \quad i = 1, \cdots, t(r).$$

(2) If r is odd,

$$\lambda_{2t(r)+1-i}(k) = -\lambda_i(k - \pi), \quad k \in [0, \pi], \quad i = 1, \cdots, t(r).$$

Proof. We first prove (i). Suppose that r is even. Because, as mentioned above, $D(\lambda)$ is an even function in this case, we have

$$D(-\lambda_i(k)) = \cos k.$$

Hence, $-\lambda_i(k)$ is also a solution of (4.1). From Lemma 4.2, we obtain the desired result.

In the case where r is odd, from an argument similar to that mentioned above, we have

$$D(-\lambda_i(k-\pi)) = \cos k.$$

This proves (ii), again by Lemma 4.2. □

Next we present the

Proof of Theorem 1.6. Because, from (4.6), we have

$$\operatorname{Arc}\cos D(\lambda_i(k)) = k, \quad k \in [0, \pi],$$

we know that $\lambda_i(k)$ $(i = 1, 2, \cdots, r + s(r) - q(r))$ are strictly monotone functions in $k \in [0, \pi]$. From Lemma 4.3, we have

$$\{\lambda_i(k) \mid k \in [0, \pi]\} = [a_i, b_i]$$

with $a_i = \min\{\lambda_i(0).\lambda_i(\pi)\}$ and $b_i = \max\{\lambda_i(0).\lambda_i(\pi)\}$. Because we know from the first half of Theorem 1.1 (2) and Lemma 4.4 that

$$\lambda_{t(r)}(k) \geq a_{t(r)} > 0 > b_{t(r)+1} \geq \lambda_{t(r)+1}(k),$$

it follows from (4.5) and Lemma 4.4 once again that

$$\sigma(L_G) \setminus \{0\} = \bigcup_{i=1}^{t(r)} [a_i, b_i] \cup [-b_i, -a_i].$$

From (2.16), we know that

$$A_{m_0} = (-1)^{t(r)}(B_{m_0} + C_{m_0}).$$

Let us first assume that $t(r)$ is even. Then, from Lemma 2.6, we have

$$\begin{cases} \lim_{\lambda \to 0} D(\lambda) = +\infty & \text{if } s(r) > q(r), \\ \lim_{\lambda \to 0} D(\lambda) > 1 & \text{if } s(r) = q(r). \end{cases}$$

From Proposition 2.2, we have $A_{t(r)+m_0} = A_{(r+s(r)-\star)/2} > 0$. Hence, we obtain

$$a_i = \begin{cases} \lambda_i(\pi), & \text{if } i = 1, 3, \cdots, t(r) - 1, \\ \lambda_i(0), & \text{if } i = 2, 4, \cdots, t(r), \end{cases}$$

$$b_i = \begin{cases} \lambda_i(0), & \text{if } i = 1, 3, \cdots, t(r) - 1, \\ \lambda_i(\pi), & \text{if } i = 2, 4, \cdots, t(r). \end{cases}$$

In the case where $t(r)$ is odd, we have

$$\begin{cases} \lim_{\lambda \to 0} D(\lambda) = -\infty & \text{if } s(r) > q(r), \\ \lim_{\lambda \to 0} D(\lambda) < -1 & \text{if} s(r) = q(r). \end{cases}$$

From an argument similar to that mentioned above, we know that

$$a_i = \begin{cases} \lambda_i(\pi), & \text{if } i = 1, 3, \cdots, t(r), \\ \lambda_i(0), & \text{if } i = 2, 4, \cdots, t(r) - 1, \end{cases}$$
$$b_i = \begin{cases} \lambda_i(0), & \text{if } i = 1, 3, \cdots, t(r), \\ \lambda_i(\pi), & \text{if } i = 2, 4, \cdots, t(r) - 1. \end{cases}$$

This completes the proof of the theorem. □

4.2 Absence of nonzero eigenvalues

We now prove part (3) of Theorem 1.1. From [5, Theorem XIII.85], we may observe that λ is an eigenvalue of L_G if and only if the Lebesgue measure of the set $\{k \in [-\pi, \pi] \mid \lambda \text{ is an eigenvalue of } L_k\}$ is positive. Hence, it suffices to show that for any $\lambda \in \sigma(L_G) \setminus \{0\}$, the Lebesgue measure of

$$E_\lambda = \{k \in [-k, k] \mid F(k, \lambda) = 0\}$$

is 0.

Let us assume that $\lambda_0 \in \sigma(L_G)$. Then we know, from (4.5), that there exists an i such that $\lambda_0 \in \{\lambda_i(k) \mid k \in [0, \pi]\}$. Because λ_i is strictly monotone in $[0, \pi]$, there exists a $k_0 \in [0, \pi]$ such that $\lambda_0 = \lambda_i(k_0)$ and $E_{\lambda_0} = \{\pm k_0\}$. Hence, the Lebesgue measure of E_{λ_0} is 0, and part (3) of Theorem 1.1 is proven.

4.3 Multiplicity

We still have to prove Lemma 4.2. From (2.8), we know that

$$\dim \ker(L_k - \lambda) = \dim \ker A_k(\lambda).$$

By definition, we have

$$A_k(\lambda) = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22}, \end{pmatrix} \tag{4.7}$$

where $A_{11} = A_{11}(k, \lambda) \in M_r(\mathbb{C})$, $A_{21} = {}^{\mathrm{t}}A_{12} \in M_{s,r}(\mathbb{C})$, $A_{22} \in M_s(\mathbb{C})$ and

$$A_{11}(k,\lambda) = \begin{pmatrix} -d_1\lambda & 1 & 0 & \cdots & 0 & e^{-ik} \\ 1 & -d_2\lambda & \ddots & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \ddots & \ddots & 1 \\ e^{ik} & 0 & \cdots & 0 & 1 & -d_r\lambda \end{pmatrix},$$

$$A_{22} = \begin{pmatrix} -\lambda & 0 & \cdots & 0 \\ 0 & -\lambda & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & -\lambda \end{pmatrix}.$$

Here $A_{12} = (e_{i,j})$ is given by

$$e_{i,j} = \begin{cases} 1, & i = i_p, j = p \\ 0, & \text{otherwise} \end{cases}.$$

From direct calculation, we have

$$A_{12}A_{21} = \mathrm{diag}(\delta_1, \cdots, \delta_r),$$

where

$$\delta_i = \begin{cases} 1, & i = i_p,\ p = 1, \cdots, s, \\ 0, & \text{otherwise.} \end{cases}$$

Henceforth, we assume that $\lambda \neq 0$. Then A_{22} has the inverse $A_{22}^{-1} = \lambda^{-1}I \in M_s(\mathbb{C})$. Hence, we obtain

$$A_{12}A_{22}^{-1}A_{21} = \lambda^{-1}\mathrm{diag}(\delta_1, \cdots, \delta_r). \tag{4.8}$$

Let $\boldsymbol{x} = {}^{\mathrm{t}}(\boldsymbol{x}_1, \boldsymbol{x}_2) \in \mathbb{C}^r \oplus \mathbb{C}^s$. From (4.7) and (4.8), we know that $A_k(\lambda)\boldsymbol{x} = 0$ if and only if

$$\begin{cases} A_{11}\boldsymbol{x}_1 = \lambda^{-1}\mathrm{diag}(\delta_1, \cdots, \delta_r)\boldsymbol{x}_1, \\ \boldsymbol{x}_2 = -A_{22}^{-1}A_{21}\boldsymbol{x}_1. \end{cases} \tag{4.9}$$

From (4.9), we observe that

$$\dim\ker(A_{11} - \lambda^{-1}\mathrm{diag}(\delta_1, \cdots, \delta_r)) = \dim\ker A_k(\lambda).$$

To prove Lemma 4.2, it therefor suffices to examine the dimension of the kernel of the matrix $E := A_{11} - \lambda^{-1}\mathrm{diag}(\delta_1, \cdots, \delta_r)$. Let $E\boldsymbol{x}_1 = 0$ and $\boldsymbol{x}_1 = {}^{\mathrm{t}}(x_1, \cdots, x_r) \in \mathbb{C}^r \setminus \{0\}$. Then, from (4.9), we have

$$x_1 = e^{-ik} g_r x_r - e^{-ik} x_{r-1} \tag{4.10}$$

$$x_2 = g_1 x_1 - e^{-ik} x_r \tag{4.11}$$

$$x_i = \lambda^{-1}(\delta_i + d_{i-1}\lambda^2) x_{i-1} - x_{i-2}, \quad 3 \le i \le r, \tag{4.12}$$

where we have set $g_i \equiv g_i(\lambda) := \lambda^{-1}(\delta_i + d_i \lambda^2)$, $i = 1, \cdots, r$. Because, from (4.11) and (4.12), we can write

$$x_i = c_i^{(1)} x_1 + c_i^{(r)} x_r, \quad i = 1, \cdots, r$$

for some $c_i^{(1)}$ and $c_i^{(r)}$, we have

$$\boldsymbol{x}_1 = x_1 \boldsymbol{c}_1 + x_r \boldsymbol{c}_r \tag{4.13}$$

with $\boldsymbol{c}_1 = {}^{\mathrm{t}}(c_1^{(1)}, \cdots, c_r^{(1)})$ and $\boldsymbol{c}_r = {}^{\mathrm{t}}(c_1^{(r)}, \cdots, c_r^{(r)})$. Hence, $\dim \ker E \le 2$ and part (2) of Lemma 4.2 is proven.

Next we prove part (1) of the lemma. Let x_i $(i = 1, \cdots, r)$ be as described above. From (4.12), we know that

$$\begin{pmatrix} x_r \\ x_{r-1} \end{pmatrix} = K \begin{pmatrix} x_2 \\ x_1 \end{pmatrix}, \tag{4.14}$$

where

$$K = \begin{pmatrix} g_{r-1} & -1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} g_2 & -1 \\ 1 & 0 \end{pmatrix}.$$

We also have

Lemma 4.5. The following holds:

$$(e^{ik} I - T) \begin{pmatrix} x_1 \\ x_r \end{pmatrix} = 0, \tag{4.15}$$

where $T = (T_{i,j})$ is given by

$$\begin{aligned}
T_{11} &= g_1(g_r K_{11} - K_{21}) + g_r K_{12} - K_{22}, \\
T_{12} &= -e^{-ik}(g_r K_{11} - K_{21}), \\
T_{21} &= e^{ik}(g_1 K_{11} + K_{12}), \\
T_{22} &= -K_{11}
\end{aligned}$$

with $K = (K_{i,j})$.

Proof. From (4.10), we have

$$x_{r-1} = g_r x_r - e^{ik} x_1. \tag{4.16}$$

Substituting (4.11) and (4.16) in (4.14), we obtain

$$\begin{pmatrix} 0 & 1 \\ -e^{ik} & g_r \end{pmatrix} \begin{pmatrix} x_1 \\ x_r \end{pmatrix} = K \begin{pmatrix} g_1 & -e^{-ik} \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_r \end{pmatrix}. \tag{4.17}$$

Several calculations establish (4.15). □

By (4.16), we have

$$(e^{ik} - T_{11})x_1 - T_{12}x_r = 0, \tag{4.18}$$

$$-T_{21}x_1 + (e^{ik} - T_{22})x_r = 0. \tag{4.19}$$

By definition, g_i $(i = 1, \cdots, r)$ are real numbers, so are $K_{i,j}$, T_{11} and T_{22}. Assume that $k \in (-\pi, 0) \cup (0, \pi)$. Then we have $(e^{ik} - T_{22}) \neq 0$ and, by (4.19),

$$x_r = \frac{T_{21}x_1}{e^{ik} - T_{22}}. \tag{4.20}$$

If $T_{21} = 0$, then $g_1 K_{11} + K_{12} = 0$. Hence, we know from (4.17) that

$$(1 + e^{-ik}K_{11})x_r = 0.$$

Because $1 + e^{-ik}K_{11} \neq 0$, we have $x_r = 0$. From (4.18) and the fact that $e^{ik} - T_{11} \neq 0$, we have $x_1 = 0$. However, this contradicts $\boldsymbol{x}_1 \neq 0$. Thus, we know that $T_{21} \neq 0$. From (4.13) and (4.20), we have $\boldsymbol{x}_1 = x_1(\boldsymbol{c}_1 + (e^{ik} - T_{22})^{-1}T_{21}\boldsymbol{c}_r)$. This completes the proof of Lemma 4.5.

5 Two dimensional lattice

In this section, we first form a class $\mathscr{G}^d$ $(d \geq 1)$ of graphs that are obtained from $\mathbb{Z}^d$ by adding pendant edges in a manner similar to the way in which $G \in \mathscr{G}$ is obtained from $\mathbb{Z}$. We will say that $G \in \mathscr{G}^d$ if G satisfies the following:

($\mathrm{G}_1^{(d)}$) $\{(\boldsymbol{n}, 0) \in \mathbb{Z}^d \times \{0, 1\} \mid \boldsymbol{n} \in \mathbb{Z}^d\} \subset V(G) \subset \mathbb{Z}^d \times \{0, 1\}$ and a vertex $(\boldsymbol{n}, 0) \in V(G)$ is connected to $(\boldsymbol{n} \pm 1, 0) \in V(G)$.

($\mathrm{G}_2^{(d)}$) If $(\boldsymbol{n}, 1) \in V(G)$, then $(\boldsymbol{n}, 1)$ is an end vertex connected to $(\boldsymbol{n}, 0)$ and $\deg(\boldsymbol{n}, 0) = 2d + 1$. If $(\boldsymbol{n}, 1) \notin V(G)$, then $\deg(\boldsymbol{n}, 0) = 2d$.

($\mathrm{G}_3^{(d)}$) There exist vectors $\boldsymbol{a}_1, \ldots, \boldsymbol{a}_d \in \mathbb{Z}^d$ linearly independent (as vectors in $\mathbb{R}^d$) such that $(\boldsymbol{n} + \boldsymbol{a}_1, 1), \ldots, (\boldsymbol{n} + \boldsymbol{a}_d, 1) \in V(G)$ if and only if $(\boldsymbol{n}, 1) \in V(G)$.

It is clear that $\mathscr{G}^1 = \mathscr{G}$ by definition.

Let $d = 2$. We consider two graphs $G_j = (V(G_j), E(G_j)) \in \mathscr{G}^2$ $(j = 1, 2)$ satisfying the condition that $(\boldsymbol{n}, 1) \in V(G_j)$ if and only if

$$\boldsymbol{n} = \sum_{i=1}^{2} n_i \boldsymbol{a}_i^{(j)}, \quad (n_1, n_2) \in \mathbb{Z}^2,$$

where $\boldsymbol{a}_1^{(1)} = (2, 0)$, $\boldsymbol{a}_2^{(1)} = (0, 1)$, $\boldsymbol{a}_1^{(1)} = (2, 0)$ and $\boldsymbol{a}_2^{(1)} = (1, 1) \in \mathbb{Z}^2$. Clearly, G_j satisfies the condition ($\mathrm{G}_3^{(d)}$) with $\boldsymbol{a}_i = \boldsymbol{a}_i^{(j)}$. The following theorem reveals that the spectrum of graphs belonging to $\mathscr{G}^2$ is more complex than the spectrum of those belonging to $\mathscr{G}^1$.

Theorem 5.1. Let $G_j \in \mathscr{G}^2$ $(j = 1, 2)$ be defined as above. Then:

(a) L_{G_1} has no spectral gap, *i.e.* $\sigma(L_{G_1}) = [-1, 1]$.

(b) L_{G_2} has a spectral gap, *i.e.* $\sigma(L_{G_2}) \neq [-1, 1]$.

Proof. By an argument similar to that in the proof of of Proposition 2.2, we observe that $\lambda \in \sigma(L_{G_1})$ if and only if there exists a pair $(k_1, k_2) \in [-\pi, \pi]^2$ such that

$$20\lambda^3 - 18(\cos k_2)\lambda^2 - 2(3 - 2\cos^2 k_2)\lambda + 2\cos k_2 - 2(\cos k_1)\lambda = 0. \quad (5.1)$$

Let $u = \cos k_1$ and $v = \cos k_2$. Then we have

$$2\lambda v^2 + (1 - 9\lambda^2)v + \lambda(10\lambda^2 - u - 3) = 0. \quad (5.2)$$

We now consider a solution of the form $\lambda = v/a$. In this case, (5.2) reduces to

$$v\{(2a - 5)(a - 2)v^2 - a^2(u + 3 - a)\} = 0 \quad (5.3)$$

Let $a = 2$. Then, from (5.3) we observe that $\lambda = v/2$ is the solution of (5.2) for any v in the range $-1 \leq v \leq 1$ if $u = -1$. Hence we have $[-1/2, 1/2] \subset \sigma(L_{G_1})$. Now let $a = 1$. It is easy to observe from (5.3) that $\lambda = v$ is the solution of (5.2) whenever $|v| = \sqrt{(u + 2)/3}$. Since $-1 \leq u \leq 1$, we have $[-1, -1/\sqrt{3}] \cup [1/\sqrt{3}, 1] \subset \sigma(L_{G_1})$. Finally, let $a = 1/2$. From (5.3) we can show that $\lambda = 2v$ is a solution of (5.3) if $|v| = \sqrt{(2u + 5)/28}$. Hence we have $[-\sqrt{7/12}, -1/2] \cup [1/2, \sqrt{7/12}] \subset \sigma(L_{G_1})$. Thus, (a) is established.

Similarly to the proof of (a), we observe that $\lambda \in \sigma(L_{G_2})$ if and only if there exists a pair $(k_1, k_2) \in [-\pi, \pi]^2$ such that

$$\lambda \left\{ 20\lambda^2 - 6 - 4\cos k_2 - 4\cos^2 k_2 - 2(1 + 2\cos k_2)\cos k_1 \right\} = 0. \tag{5.4}$$

We shall prove that $\lambda \in \rho(L_{G_1})$ if $0 < \lambda < 1/\sqrt{5}$. We fix $\lambda_0 \in (0, 1/\sqrt{5})$ and assume that $\lambda_0 \in \sigma(L_{G_1})$. Let $u = \cos k_1$ and $v = \cos k_2$. Then we know, from (5.4), that there exist a pair $(u, v) \in [-1, 1]^2$ such that

$$v^2 + (1 + u)v + \frac{3 + u - 10\lambda_0^2}{2} = 0.$$

Because the above quadratic polynomial in v has a solution, its discriminant satisfies $u^2 + 20\lambda_0^2 - 5 \geq 0$. On the other hand, by assumption, we have $u^2 + 20\lambda_0^2 - 5 < 0$. This is a contradiction, and (b) is therefore proven. ☐

Acknowledgments: The author would like to thank Daiki Hayashi for useful discussions. This work was supported by Grant-in-Aid for Young Scientists(Start-up) 22840022.

References

[1] P. Chandrachud, B. S. Pujari, S. Haldar, B. Sanyal and D. G. Kanhere, A Systematic Study of Electronic Structure from Graphene to Graphane, *J. Phys.: Condens. Matter* **22** (2010), 465502.

[2] Y. Higuchi and Y. Nomura, Spectral structure of the Laplacian on a covering graph, *European J. Combin.* **30** (2009), 570 – 585.

[3] T. Kato, *Perturbation Theory for Linear Operators*, 2nd Edition, Springer, Berlin Heidelberg New York (1976).

[4] A. Suzuki, Spectrum of the Laplacian on a covering graph with pendant edges II: lattices in two dimensions, in preparation.

[5] M. Reed and B. Simon, *Methods of Modern Mathematical Physics Vol. IV*, Academic Press, New York, 1978.